



1811 2<sup>nd</sup> St SW  
Rochester, MN  
289-6068

7955 18<sup>th</sup> Ave NW  
Rochester, MN  
289-0022

[www.sargentsgardens.com](http://www.sargentsgardens.com)

## **Pruning Clematis**

Clematis can be a confusing group of plants to prune, since they are not all pruned the same way. There are three methods that can be applied to major groups depending on the time of year the plant flowers. The earliest flowering clematis bloom on old wood, while later flowering types must produce new growth in order for flower buds to form. Prune carefully, since vines are usually well entangled.

### **Group A: Early-flowering Clematis**

Plants in this group bloom in early spring, generally in April and May, from buds produced the previous season. Prune these plants immediately after flowering, but no later than the end of July. This allows time for new growth to produce flower buds for the next season. Remove shoots that have bloomed. You can prune out more vines to reduce the size or to form a good framework of branches, but avoid cutting into woody trunks.

#### **Plants in this group include:**

C. alpine, C. macropetala, C. armandii, C. Montana, C. chrysocoma and Stolwijk Gold

### **Group B: Large-flowered Hybrids**

Large-flowered hybrids bloom in mid-June on short stems from the previous season's growth and often again in late summer on new growth, though these blooms are usually smaller. Prune in February or March by removing dead and weak stems, then cut back the remaining stems to the topmost pair of large, plum green buds. This cut could be 6-18" from the stem tips. Plants in this group have the tendency to become leafless at the base as they mature. You can under plant with low, spreading perennials to help conceal the stems. You may be able to force a flush of new growth from the base by cutting the vine back to 18" immediately after the flush of bloom in June.

#### **Plants in this group include:**

The President, Integrifolia, Bourbon, Carnaby, Claire de Lune, Vino, Diamantina, Shimmer, Belle of Woking, Charissima, Crystal Fountain, Dr Ruppel, Duchess of Edinburgh, Elsa Spaeth, Fireworks, Gillian Blades, Guernsey Cream, Henryi, Josephine, Killian Donahue, Multi-Blue, Nelly Moser, Niobe, Perrins Pride, Ramona, Rebecca and Royalty.

### **Group C: Late-flowering Clematis**

Plants in this group flower on the last 24-36 inches of the current season's growth. Some types begin blooming in mid-June and continue into the fall. This is the easiest group to prune since no old wood needs to be maintained. In February or March cut each stem to a height of about 24-36 inches. This will include removal of some good stems and buds. Eventually the length of the bare stem at the base will increase as the vine matures.

#### **Plants in this group include:**

Comtesse de Bouchard, Hagley Hybrid, Jackmanii, Jackmanii Superbra, Mime. Baron Veillard,  
Prince Charles, Rhapsody, Polish Spirit, Venosa Violacea, Paniculata, Roguchi, Tangutica,  
Duchess of Albany, Princess Diana, Etoile Violette, Rosemoor,